

ERDC/EL TR-01-34

Environmental Laboratory



**US Army Corps
of Engineers®**
Engineer Research and
Development Center

Macroinvertebrate Communities and Habitat in Luxapalila Creek, Mississippi and Alabama

Barry S. Payne, Peter Smiley, and Andrew C. Miller

September 2001

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.



PRINTED ON RECYCLED PAPER

Macroinvertebrate Communities and Habitat in Luxapalila Creek, Mississippi and Alabama

by Barry S. Payne , Andrew C. Miller

Environmental Laboratory
U.S. Army Engineer Research and Development Center
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Peter Smiley

Mississippi State University
Mississippi State, MS 39762

Final report

Approved for public release; distribution is unlimited

Contents

Preface	vi
1—Introduction	1
2—Study Area and Methods.....	2
Study Area.....	2
Site Selection and Data Collection	2
Before and After Project Comparisons.....	4
Data Analysis	4
3—Results and Discussion	5
Habitat Characteristics	5
Macroinvertebrate Community Characteristics	7
Before and After Project Comparisons.....	11
4—Summary.....	22
References	24
SF 298	

List of Figures

Figure 1.	Study sites of Luxapalila Creek, Mississippi and Alabama.....	3
Figure 2.	Relative abundance of functional feeding groups of macroinvertebrates from fall samples from pools in three channel types	12
Figure 3.	Relative abundance of functional feeding groups of macroinvertebrates from spring samples from pools in three channel types	13

Figure 4.	Relative abundance of functional feeding groups of macroinvertebrates from fall samples from riffles in three channel types	14
Figure 5.	Relative abundance of functional feeding groups of macroinvertebrates from spring samples from riffles in three channel types	15

List of Tables

Table 1.	Summary of Mean Geomorphological and Vegetative Characteristics of Streambanks Within the Recently Channelized, Historically Channelized, and Unchannelized Portions of Luxapalila Creek	5
Table 2.	Summary of Mean Habitat Parameters of Pools Among All Channel Types and Sampling Periods	6
Table 3.	Summary of Mean Habitat Parameters of Riffles Among All Channel Types and Sampling Periods	7
Table 4.	Summary of Two Factor ANOVA Results and Factor Means of Pool Habitat Parameters That Exhibited Significant Single Factor Effects.	8
Table 5.	Summary of Two Factor ANOVA Results and Factor Means of Riffle Habitat Parameters That Exhibited Significant Single Factor Effects	9
Table 6.	Relative Abundance of Macroinvertebrate Taxa From Pools in Three Channel Types and Two Sampling Periods.....	10
Table 7.	Relative Abundance of Macroinvertebrate Functional Groups From Pools in Three Channel Types and Two Sampling Periods.....	11
Table 8.	Relative Abundance of Macroinvertebrate Functional Groups From Riffles in Three Channel Types and Two Sampling Periods.....	11
Table 9.	Summary of Two Factor ANOVA Results and Factor Means of the Density of the Five Most Common Macroinvertebrate Taxa in Pools That Exhibited Significant Single Factor Effects	16

Table 10.	Summary of Two Factor ANOVA Results and Factor Means of the Density of the Five Most Common Macroinvertebrate Taxa in Riffles That Exhibited Significant Single Factor Effects	16
Table 11.	Mean Density of Macroinvertebrate Taxa From Pools in Three Channel Types and Two Sampling Periods	17
Table 12.	Mean Density of Macroinvertebrate Taxa From Riffles in Three Channel Types and Two Sampling Periods	19
Table 13.	Relative Abundance of Macroinvertebrate Taxa From Pools in the Recently Altered Portion of Luxapalila Creek Among Pre- and Post-Alteration Sampling Periods and Two Seasons	20
Table 14.	Relative Abundance of Macroinvertebrate Taxa From Riffles in the Recently Altered Portion of Luxapalila Creek Among Pre- and Post-Alteration Sampling Periods and Two Seasons	21

Preface

The study reported herein was conducted by the U.S. Army Engineer Research and Development Center (ERDC), Vicksburg, MS, for the U.S. Army Engineer District, Mobile. The purpose was to analyze macroinvertebrate community composition and habitat at a series of pools and riffles in Luxapalila Creek in relation to recent construction of flood control measures in the lower portion of that stream.

Fieldwork was conducted by Mr. Peter Smiley, Mississippi State University (MSU), with the assistance of Mr. Will Green, University of Southern Mississippi. Drs. Andrew C. Miller, Environmental Laboratory (EL), ERDC, and Eric Dibble, MSU, provided guidance during conduct of this study.

During the conduct of this study, Dr. Edwin A. Theriot was Acting Director, EL; Dr. Dave Tazik was Chief, Ecosystem Evaluation and Engineering Division (EEED), EL; and Mr. Larry Sanders was Acting Chief, Aquatic Ecology and Invasive Species Branch (AEIB), EEED. Authors of this report were Drs. Barry S. Payne and Andrew C. Miller, AEIB, and Mr. Smiley.

At the time of publication of this report, Director of ERDC was Dr. James R. Houston. Commander and Executive Director was COL John W. Morris III, EN.

This report should be cited as follows:

Payne, B. S., Smiley, P., and Miller, A. C. (2001). "Macroinvertebrate communities and habitat in Luxapalila Creek, Mississippi and Alabama," ERDC/EL TR-01-34, U.S. Army Engineer Research and Development Center, Vicksburg, MS.

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

1 Introduction

The U.S. Army Engineer District, Mobile constructed flood control measures in the lower reach of Luxapalila Creek from 1998-2000. These measures included clearing and snagging, bank protection, excavation of pools, and construction of rock rubble weirs. The measures were designed to protect rural areas upstream of Columbus, MS, against floods expected to occur every 1.5 years, and to protect urban areas in Columbus from floods expected to occur every 5 years. Many aquatic habitats are altered by channel diversion, modification, or construction of dams (Standford and Ward 1979). Consequently, potentially adverse impacts to stream biota are a concern.

From 1987-1989, a study was conducted of stream characteristics and the macroinvertebrate community at a set of four riffles and four pools in an approximately 30-mile (48-km) reach of Luxapalila Creek extending from Columbus, MS, to Millport, AL (Payne et al. 1991; Payne and Miller 1991). The purpose of that study was to describe baseline conditions that could be subsequently compared to conditions revealed by a similar post-project study. This report presents the results of the post-project study of physical and biological conditions in Luxapalila Creek.

Emphasis in both studies was on macroinvertebrates, especially the more sessile taxa of fully aquatic organisms and aquatic life stages of insects. These animals are often used to monitor stream conditions and changes (Barbour et al. 1999). Density and diversity of these animals reflect geomorphologic, substratum, and water quality conditions.

2 Study Area and Methods

Study Area

Luxapalila Creek watershed is situated within the northern edge of the Gulf Coastal Plains Physiographic Province. Luxapalila Creek is a tributary of the Tombigbee River and originates in Alabama and flows through northeast Mississippi (Payne et al. 1991). A 41.8-km section of Luxipalila Creek extending from Winfield, AL, to the Alabama-Mississippi state line was channelized in 1922, and then the lower 3.4 km was channelized in 1967 (Payne et al. 1991). In 1996, channelization and widening of the lower reaches near Columbus, MS, occurred.

Currently, three channel types are present within Luxapalila Creek (Figure 1). The downstream portion includes a recently channelized reach near Columbus, MS. The middle reach, near Steens, MS, has never been channelized. The very straight, upstream reach in Alabama was channelized approximately 80 years ago.

Site Selection and Data Collection

Two study sites were selected from within each of the different channel types and sampled as part of the present study (Figure 1). Sites 1 and 2 are located within the recently channelized portion and correspond with sites 1 and 2 studied by Payne et al. (1991). Site 1 is located downstream of Highway 182, and site 2 is located near the confluence of Magby Creek. Sites 3 and 4 are located within the unchannelized portion of Luxapalila Creek. Site 3 is located downstream of Mississippi Highway 12, and site 4 is located downstream of Gunshoot Road. Sites 5 and 6 are located within the historically channelized portion of Luxapalila Creek in Alabama. Site 5 is near Highway 17 in Millport, AL, while site 6 is downstream of Highway 49 near Kennedy, AL.

A previous study compared the macroinvertebrate communities of pools and riffles within Luxapalila Creek (Payne et al. 1991; Payne and Miller 1991). In the present study a space for time assessment was conducted to examine how macroinvertebrate communities and physical habitat within a microhabitat type varied among the three channel types and two seasonal periods (fall and spring).




Figure 1. Study sites of Luxapalila Creek, Mississippi and Alabama

Preliminary observations indicated that distinct microhabitat types were present within the unchannelized and historically channelized portions of Luxapalila Creek. However, distinct pools and riffles were not present within the recently channelized portion of Luxapalila Creek. Pools and riffles within the recently channelized portion of Luxapalila Creek were delineated using qualitative habitat characteristics to define each microhabitat type to facilitate comparisons among channel types. Pools were considered to be areas of slow moving water greater than 0.56 m in depth with little surface turbulence, whereas riffles were defined as areas of fast moving water less than 0.56 m deep and exhibiting greater surface turbulence than pools.

Macroinvertebrates and physical habitat data were collected from each of the selected pools and riffles during four sampling periods: fall 1998, spring 1999, fall 1999, and spring 2000). Macroinvertebrates were sampled with a hand-held coring device (Miller and Bingham 1987) that sampled 0.0079 sq m. Five cores for macroinvertebrate samples were taken within each microhabitat type (total cores per sampling period = 180).

Water quality parameters (pH, temperature, dissolved oxygen, and turbidity) and estimated percent canopy cover in the middle of the sampling area of each microhabitat were measured during each sampling period. Depth, velocity, and substrate types were measured at three points along a transect located in the middle of the sampling area within each microhabitat type. Substrate types were visually classified in the field as either clay, sand, gravel, woody debris, or riprap.

Macroinvertebrate samples were preserved with 10-percent formalin solution in the field. In the laboratory, an elutriation process (Payne et al. 1991) was used to remove macroinvertebrates from the sediment. Macroinvertebrate samples were

swirled in a 12-L bucket and poured through a 250- μ mesh sieve. Each sample was elutriated five times. Previous work indicated that this process removes 90 to 100 percent of macroinvertebrates within a sample (Payne et al. 1991). Macroinvertebrates were removed from elutriated samples with the aid of a stereomicroscope. Organisms were first sorted to major groups, such as chironomids, oligochaetes, and ephemeropterans, and counted.

Following the initial identifications, all sorted samples were recounted and further identifications were made. The following animals were only identified to major group (typically class): segmented worms (Oligochaetes), roundworms (Nematodes), horsehair worms (Nematomorpha), water mites (Hydracarina), planaria (Platyhelminthes), leeches (Hirudinea), amphipods (Amphipoda), crayfish (Decapoda), isopods (Isopoda), snails (Gastropoda), mussels (Bivalvia), springtails (Collembola), crickets (Orthoptera), and megalopterans (Megaloptera). The following taxa were identified to family level when possible: Coleoptera, Diptera, Ephemeroptera, Hemiptera, Odonata, Plecoptera, and Trichoptera. This level of taxonomic resolution allowed enumeration per functional feeding group.

Before and After Project Comparisons

Sites 1 and 2 (of the present study) were located within the recently channelized reach that was represented by sites 1, 2, and 3 of the previous study (Payne et al. 1991). Sites 5 and 6 (of the present study) fell in the historically channelized reach, as did site 4 of the previous study. At a level of river reach, comparisons were thus possible of pre- and post-project conditions in the recently channelized and historically channelized portions of the stream. At a finer level of spatial detail, riffles and pools from sites 1 and 2 of Payne et al. (1991) were within 200 m of sites 1 and 2 of the present study. Therefore, macroinvertebrate community structure at those two sites was specifically compared before and after project construction. Sampling methods and efforts with respect to macroinvertebrates were equal between pre- and post-alteration periods. In the 1981-1989 study, taxonomic resolution involved genus- and species-level identification of oligochaetes and dipterans to the maximum extent feasible. In the present study, identifications were made to the family level.

Data Analysis

Differences in mean physical habitat parameters among the different channel types and seasonal periods were assessed using a two factor analysis of variance (ANOVA) coupled with Student Newman-Keul's SNK multiple range test. Assumptions of ANOVA were tested prior to statistical testing, and if the data did not meet the assumptions, then they were transformed using $\log(x + 1)$ transformation. Variables calculated as percents (percent canopy cover, percent sand, etc.) were arcsine square root transformed prior to statistical testing (Zar 1987). Univariate statistical tests were conducted using SigmaStat 2.0 for Windows (Jandel Corporation 1995). Significance level for all univariate tests was $P < 0.05$.

3 Results and Discussion

Habitat Characteristics

Geomorphological and vegetative characteristics of streambanks differed among recently channelized (RC), historically channelized (HC), and unchannelized (UC) portions of Luxapalila Creek (Table 1). Banks were steeper in the HC and UC than RC portions of the creek. The stream was narrowest in the UC portion. Trees were almost absent and shrubs were not common in the RC portion. Riprap was abundant in the RC but not the other portions. Bare soil comprised half of the streambank in the UC portion, one fifth in the RC portion, but one tenth in the HC portion of the creek. Canopy cover was absent in the RC portion of the creek, high to moderate in the HC portion, and moderate in the UC portion.

Characteristic	RC	HC	UC
Bank angle, deg	17.63	29.36	27.87
Bank height, m	5.15	3.52	1.98
Top bank width, m	57.54	42.06	26.93
Bottom channel width, m	28.21	29.12	17.84
Percent trees (woody vegetation > 5 m)	0.03	5.35	6.25
Percent shrub (woody vegetation < 5 m)	11.71	34.38	22.50
Percent herbaceous vegetation	39.05	49.72	23.82
Percent riprap	27.89	2.99	0.42
Percent bare soil	21.32	7.71	47.01

Despite geomorphic and vegetative cover differences, water quality in pools (Table 2) and riffles (Table 3) was good and similar among all three portions of the creek. Measurements of pH indicated a very slightly acidic condition (6.7 to 6.8). Dissolved oxygen was high (9.7 to 10.4 mg/L) at temperatures ranging from 12.5 to 15.7 °C. Turbidity was low (8.0 to 11.5 NTU).

Table 2 Summary of Mean (Standard Error) Habitat Parameters of Pools Among All Channel Types and Sampling Periods			
Habitat Parameter	RC	HC	UC
Fall			
pH	6.77 (0.20)	6.84 (0.20)	6.74 (0.17)
Temperature, °C	15.60 (1.22)	12.40 (1.15)	14.20 (1.16)
Dissolved oxygen, mg/L	9.98 (0.18)	10.41 (0.24)	9.66 (0.26)
Turbidity, NTU	8.44 (2.95)	8.04 (2.95)	11.46 (4.19)
Depth, m	0.64 (0.06)	0.72 (0.08)	0.71 (0.06)
Velocity, m/sec	0.37 (0.13)	0.08 (0.05)	0.00 (0.00)
Percent clay	25.00 (15.96)	0.00 (0.00)	0.00 (0.00)
Percent sand	0.00 (0.00)	83.33 (9.62)	25.00 (15.96)
Percent gravel	66.67 (13.61)	16.67 (9.62)	75.00 (15.96)
Percent woody debris	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent riprap	8.33 (8.33)	0.00 (0.00)	0.00 (0.00)
Percent canopy cover	0.00 (0.00)	75.00 (14.43)	25.00 (14.43)
Spring			
pH	6.61 (0.14)	7.24 (0.19)	7.00 (0.11)
Temperature, °C	23.48 (0.36)	22.23 (0.43)	23.33 (0.25)
Dissolved oxygen, mg/L	8.20 (0.16)	8.35 (0.28)	7.52 (0.13)
Turbidity, NTU	12.99 (5.78)	9.69 (3.27)	14.91 (6.72)
Depth, m	0.68 (0.06)	0.89 (0.04)	0.72 (0.03)
Velocity, m/sec	0.47 (0.19)	0.10 (0.05)	0.01 (0.01)
Percent clay	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent sand	16.67 (16.67)	58.33 (20.97)	58.33 (20.97)
Percent gravel	83.33 (16.67)	16.67 (16.67)	41.67(20.97)
Percent woody debris	0.00 (0.00)	25.00 (8.00)	0.00 (0.00)
Percent riprap	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent canopy cover	0.00 (0.00)	87.50 (12.50)	25.00 (14.43)

Pools ranged in depth from 0.64 to 0.72 m (Table 2). Velocity in the pools was very low in the HC and UC portions (0.0 to 0.1 m/sec) and moderate (0.4 m/sec) in the RC portion. Riffles ranged only from 0.28 to 0.32 m deep (Table 3); velocity was higher (0.8 m/sec) in the UC than HC (0.5 m/sec) or RC (0.4 m/sec) portions of the creek.

Substratum in pools differed markedly among the reaches (Table 2). Gravel (67 percent), hardpan clay (25 percent), and riprap (8 percent) were the only substratum types encountered in pools in the RC portion. In contrast, sand (83 percent) dominated the pools in the HC portion, while gravel (75 percent) was the predominant substratum in pools in the UC portion. Gravel was the predominant substratum (92 to 100 percent) of riffles in all stream reaches (Table 3).

ANOVA summaries of physical habitat and water quality characteristics of pools and riffles are provided in Tables 4 and 5, respectively.

Table 3 Summary of Mean (Standard Error) Habitat Parameters of Riffles Among All Channel Types and Sampling Periods			
Habitat Parameter	RC	HC	UC
Fall			
pH	6.78 (0.20)	6.83 (0.17)	6.78 (0.20)
Temperature, °C	15.70 (1.32)	12.45 (1.14)	14.18 (1.14)
Dissolved oxygen, mg/L	9.87 (0.22)	10.16 (0.19)	9.57 (0.30)
Turbidity, NTU	7.79 (3.01)	8.12 (3.13)	11.00 (3.03)
Depth, m	0.28 (0.04)	0.25 (0.06)	0.32 (0.03)
Velocity, m/sec	0.36 (0.14)	0.45 (0.09)	0.79 (0.04)
Percent clay	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent sand	0.00 (0.00)	0.08 (0.08)	0.00 (0.00)
Percent gravel	100.00 (0.00)	91.67 (8.33)	100.00 (0.00)
Percent woody debris	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent riprap	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent canopy cover	0.00 (0.00)	50.00 (28.87)	37.50 (23.94)
Spring			
pH	6.70 (0.20)	7.23 (0.18)	7.02 (0.09)
Temperature, °C	23.58 (0.51)	22.18 (0.39)	23.40 (0.25)
Dissolved oxygen, mg/L	8.13 (0.20)	8.51 (0.26)	7.78 (0.18)
Turbidity, NTU	13.56 (6.05)	8.56 (3.82)	13.33 (5.98)
Depth, m	0.45 (0.04)	0.24 (0.05)	0.52 (0.04)
Velocity, m/sec	0.67 (0.10)	0.42 (0.06)	0.87 (0.13)
Percent clay	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent sand	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent gravel	100.00 (0.00)	100.00 (0.00)	100.00 (0.00)
Percent woody debris	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent riprap	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Percent canopy cover	0.00 (0.00)	50.00 (28.87)	37.50 (23.94)

Macroinvertebrate Community Characteristics

A functional analysis of macroinvertebrate community composition indicated a remarkable degree of uniformity within the stream, regardless of season, reach, or habitat type within a reach (Tables 6-8; Figures 2-5). The macroinvertebrate community of Luxapalila Creek was consistently dominated by collector-gatherers and predators. This uniformity was true for recently channelized, historically channelized, and unchannelized stream reaches and for pools and riffles. Collector-gatherers comprised 55 to 83 percent of the fauna considering all locations and both seasons. Predatory macroinvertebrates comprised 15 to 45 percent of the fauna for both seasons. Shredders and scrapers comprised only 0 to 4 percent of the fauna, probably reflecting the lack of large gravel and cobble in the streambed for both seasons. Collector-filterers comprised from 0 to 18 percent of the fauna and were consistently least abundant in the historically channelized

Table 4 Summary of Two Factor ANOVA Results and Factor Means (Standard Error) of Pool Habitat Parameters That Exhibited Significant Single Factor Effects				
Channel Type	ANOVA Results	Factor Means		
		RC	HC	UC
pH	NS	-	-	-
Temperature, °C	NS	-	-	-
Dissolved oxygen, mg/L	S	9.09 (0.35) a	9.38 (0.43) a	8.59 (0.43) b
Turbidity, NTU	NS	-	-	-
Depth, m	S	0.66 (0.04) a	0.81 (0.05) b	0.72 (0.03) ab
Velocity, m/sec	S	0.42 (0.11) a	0.09 (0.03) b	0.004 (0.003) b
Percent clay	NS	-	-	-
Percent sand	S	8.33 (8.33) a	70.83 (11.68) b	41.67 (13.73) b
Percent gravel	S	75.00(10.45) a	16.67 (8.91) b	58.33 (13.73) ab
Percent woody debris	CT X S	-	-	-
Percent riprap	NS	-	-	-
Percent canopy cover	S	0.00 (0.00) a	81.25 (9.15) b	25.00(9.45) c
Season	ANOVA Results	Factor Means		
		Fall	Spring	
pH	NS	-	-	
Temperature, °C	S	14.07 (0.73)	23.01 (0.25)	
Dissolved oxygen, mg/L	S	10.02 (0.15)	8.02 (0.15)	
Turbidity, NTU	NS	-	-	
Depth, m	NS	-	-	
Velocity, m/sec	NS	-	-	
Percent clay	NS	-	-	
Percent sand	NS	-	-	
Percent gravel	NS	-	-	
Percent woody debris	CT X S	-	-	
Percent riprap	NS	-	-	
Percent canopy cover	NS	-	-	
Note: NS = not significant, S = significant, CT X S = significant interaction effect. Means with the same letter are not significantly different.				

upper reach of the stream (Tables 7 and 8). Collector-filterers were mostly the Asian clam, *Corbicula fluminea*, and a few native unionid mussels. These bivalves were enormously greater in size than the numerically dominant chironomids, oligochaetes, and similarly small organisms. Thus, the ecological significance of filtering to stream trophic dynamics almost certainly was greater than indicated by their numerical relative abundance.

Total density of four of the five most common macroinvertebrates in pools (midges, unsegmented worms, segmented worms, and mites) did not vary significantly among recently channelized, historically channelized, and

Table 5 Summary of Two Factor ANOVA Results and Factor Means (Standard Error) of Riffle Habitat Parameters That Exhibited Significant Single Factor Effects				
Channel type	ANOVA Results	Factor Means		
		RC	HC	UC
pH	NS	-	-	-
Temperature, °C	NS	-	-	-
Dissolved oxygen, mg/L	S	9.00 (0.36) ab	9.34 (0.35) a	8.68 (0.37) b
Turbidity, NTU	NS	-	-	-
Depth, m	S	0.36 (0.04) a	0.24 (0.04) b	0.42 (0.05) a
Velocity, m/sec	S	0.51 (0.10) a	0.43 (0.05) a	0.83 (0.06) b
Percent clay	NS	-	-	-
Percent sand	NS	-	-	-
Percent gravel	NS	-	-	-
Percent woody debris	NS	-	-	-
Percent riprap	NS	-	-	-
Percent canopy cover	NS	-	-	-
Season	ANOVA Results	Factor Means		
		Fall	Spring	
pH	NS	-	-	
Temperature, °C	S	14.11 (0.74)	23.05 (0.28)	
Dissolved oxygen, mg/L	S	9.87 (0.15)	8.14 (0.14)	
Turbidity, NTU	NS	-	-	
Depth, m	S	0.28 (0.03)	0.40 (0.04)	
Velocity, m/sec	NS	-	-	
Percent clay	NS	-	-	
Percent sand	NS	-	-	
Percent gravel	NS	-	-	
Percent woody debris	NS	-	-	
Percent riprap	NS	-	-	
Percent canopy cover	NS	-	-	
Note: NS = not significant, S = significant. Means with the same letter are not significantly different.				

unchannelized sections of the creek (Table 9). In contrast, bivalves (mostly *C. fluminea*) were much more abundant in recently channelized than historically channelized sections. Density was intermediate in the unchannelized portion of the creek. This pattern follows an upstream to downstream increase in *C. fluminea* density that might be expected due to trophic considerations, rather than channel modifications, applicable to a stream this size.

Seasonal differences in density of the most abundant taxa in pools were more evident than habitat differences (Table 9). Chironomid abundance in fall was more than twice that observed in spring. Oligochaetes were nearly four times more abundant in fall than spring. Bivalves were approximately 50 times more

Table 6
Relative Abundance (Percent) of Macroinvertebrate Taxa From Pools
in Three Channel Types and Two Sampling Periods

Order	Family	Fall			Spring		
		RC	HC	UC	RC	HC	UC
Bivalvia		6.768	0.464	1.474	0.205	0.211	0.124
Coleoptera	Dryopidae	0.000	0.000	0.200	0.000	0.000	0.000
	Dytiscidae	0.000	0.000	0.050	0.000	0.000	0.000
	Elmidae	0.400	1.404	0.914	0.328	0.136	1.068
	Gyrinidae	0.000	0.000	0.000	0.000	0.000	0.000
	Staphylinidae	0.000	0.000	0.050	0.000	0.000	0.000
	Unknown	0.000	0.000	0.000	0.000	0.093	0.000
Collembola		0.054	0.155	0.000	0.109	0.093	0.000
Diptera	Ceratopogonidae	0.749	5.032	3.445	1.218	2.270	2.009
	Chironomidae	53.382	59.990	65.658	69.097	67.269	47.313
	Empididae	0.098	0.108	0.249	0.000	0.085	0.281
	Nymphomyiidae	0.000	0.000	0.000	0.000	0.291	0.000
	Simuliidae	0.000	0.000	0.000	1.040	0.366	0.157
	Tabanidae	0.000	0.000	0.000	0.000	0.000	0.048
	Tanyderidae	0.000	0.000	0.000	0.000	0.000	0.000
	Tipulidae	0.000	0.000	0.000	0.000	0.000	0.000
	Unknown	0.366	0.000	0.100	0.041	0.117	0.051
	Ephemeroptera	Baetidae	0.000	0.000	0.000	0.000	0.175
Baetiscidae		0.000	0.000	0.125	0.000	0.000	0.000
Caenidae		0.000	0.000	0.000	0.547	0.000	0.102
Ephemeridae		0.161	0.668	0.226	0.000	0.210	0.000
Heptageniidae		1.020	0.077	0.449	0.323	0.000	0.000
Isonychiidae		0.161	0.000	0.000	0.000	0.000	0.000
Polmitarcyidae		0.000	0.000	0.000	0.000	0.000	0.000
Potamanthidae		0.000	0.000	0.000	0.164	0.093	0.000
Tricorythidae		0.000	0.077	0.000	0.657	0.000	0.191
Unknown	1.174	1.558	0.100	3.255	1.107	0.673	
Gastropoda		2.631	0.185	0.848	0.123	0.000	0.000
Hirudinea		0.166	0.000	0.000	0.000	0.000	0.000
Hydracarina		2.687	4.504	0.984	1.133	3.908	1.372
Isopoda		0.000	0.000	0.000	0.000	0.155	0.000
Megaloptera		0.000	0.000	0.000	0.000	0.233	0.000
Nematoda		14.136	13.908	18.853	12.417	15.154	40.410
Nematomorpha		0.000	0.000	0.050	0.000	0.000	0.000
Odonata	Coenagrionidae	0.128	0.000	0.000	0.000	0.000	0.153
	Corduliidae	0.000	0.309	0.000	0.000	0.000	0.000
	Gomphidae	0.054	0.103	0.266	0.000	0.093	0.000
	Unknown	0.036	0.077	0.116	0.041	0.000	0.000
Oligochaeta		13.601	7.307	4.288	8.141	6.921	5.078
Orthoptera	Tridactylidae	0.000	0.000	0.025	0.000	0.233	0.000
Plecoptera	Perlidae	0.357	0.155	0.000	0.370	0.000	0.000
	Perlodidae	0.000	0.000	0.050	0.312	0.214	0.000
	Unknown	0.000	0.116	0.000	0.000	0.133	0.000
Trichoptera	Brachycentridae	0.000	0.000	0.000	0.000	0.000	0.096
	Hydropsychidae	0.268	0.000	0.000	0.000	0.000	0.000
	Hydroptilidae	0.514	0.147	0.299	0.000	0.000	0.191
	Leptoceridae	0.612	3.619	0.831	0.000	0.000	0.281
	Philopotamidae	0.223	0.000	0.050	0.066	0.000	0.000
	Polycentropodidae	0.027	0.000	0.100	0.000	0.000	0.000
	Psychomyiidae	0.000	0.000	0.000	0.000	0.058	0.000
	Unknown	0.232	0.039	0.200	0.411	0.383	0.306

Table 7 Relative Abundance (Percent) of Macroinvertebrate Functional Groups From Pools in Three Channel Types and Two Sampling Periods						
Invertebrate Group	Fall			Spring		
	RC	HC	UC	RC	HC	UC
Collector-filterer	7.49	0.46	1.63	1.48	0.67	0.38
Collector-gatherer	69.80	74.81	72.36	82.51	76.51	55.00
Scraper	3.67	0.26	1.30	0.45	0.29	0.00
Shredder	0.00	0.00	0.23	0.00	0.23	0.00
Predator	18.52	24.32	24.19	15.56	22.29	44.43
Piercer	0.52	0.15	0.30	0.00	0.00	0.19

Table 8 Relative Abundance (Percent) of Macroinvertebrate Functional Groups From Riffles in Three Channel Types and Two Sampling Periods						
Invertebrate Group	Fall			Spring		
	RC	HC	UC	RC	HC	UC
Collector-filterer	3.94	2.11	6.39	1.21	0.67	18.18
Collector-gatherer	71.73	79.80	62.07	83.03	81.93	63.32
Scraper	4.18	0.63	2.12	0.80	0.00	0.78
Shredder	0.00	0.19	0.00	0.00	0.21	0.00
Predator	19.41	16.95	27.65	14.90	17.06	17.55
Piercer	0.74	0.32	1.77	0.06	0.14	0.17

abundant in fall than spring. However, nematodes and mites did not show such seasonal differences.

In riffles, total density of chironomids, oligochaetes, and nematodes varied significantly among recently channelized, historically channelized, and unchannelized portions of the stream (Table 10). All taxa were highest in density in the recently channelized reach and least dense in the historically channelized reach. Density tended to be intermediate in the unchannelized reach. As with *C. fluminea*, this patterns follows an upstream to downstream increase in density. Seasonal difference in macroinvertebrate density in riffles was similar to that seen in pools. Chironomids, oligochaetes, and nematodes were all denser in fall than spring (Table 10).

Before and After Project Comparisons

At the level of river reach, total macroinvertebrate density, dominated by chironomids, was substantially greater in the present study (Tables 11 and 12) than in the 1987-1989 study (Figure 3 in Payne and Miller 1991). In the previous study, a pool in the historically channelized reach supported 6,000 to




Figure 2. Relative abundance of functional feeding groups of macroinvertebrates from fall samples from pools in three channel types




Figure 3. Relative abundance of functional feeding groups of macroinvertebrates from spring samples from pools in three channel types




Figure 4. Relative abundance of functional feeding groups of macroinvertebrates from fall samples from riffles in three channel types




Figure 5. Relative abundance of functional feeding groups of macroinvertebrates from spring samples from riffles in three channel types

Table 9
Summary of Two Factor ANOVA Results and Factor Means
(Standard Error) of the Density of the Five Most Common
Macroinvertebrate Taxa in Pools That Exhibited Significant Single
Factor Effects

Channel type	ANOVA Results	Factor Means		
		RC	HC	UC
Chironomidae	NS	-	-	-
Nematoda	NS	-	-	-
Oligochaeta	NS	-	-	-
Bivalvia	S	1616.95 (870.47) a	87.40 (61.25) b	195.29 (78.43) ab
Hydracarina	NS	-	-	-

Season	ANOVA Results	Factor Means	
		Fall	Spring
Chironomidae	S	20525.60 (1615.15)	8073.17 (1852.82)
Nematoda	NS	-	-
Oligochaeta	S	3305.81 (1169.28)	892.92 (246.67)
Bivalvia	S	1242.76 (588.54)	23.67 (6.64)
Hydracarina	NS	-	-

Note: NS = not significant, S = significant. Means with the same letter are not significantly different.

Table 10
Summary of Two Factor ANOVA Results and Factor Means
(Standard Error) of the Density of the Five Most Common
Macroinvertebrate Taxa in Riffles That Exhibited Significant Single
Factor Effects

Channel type	ANOVA Results	Factor Means		
		RC	HC	UC
Chironomidae	S	27287.21 (5033.57) a	14040.72 (3916.88) b	17425.80 (3221.34) ab
Oligochaeta	S	6275.66 (2171.57) a	1140.99 (447.65) b	3334.33 (759.89) a
Nematoda	S	6110.52 (1803.20) a	993.16 (326.39) b	3588.065 (988.61) a
Ceratapogonidae	NS	-	-	-
Ephemeroptera (unknown)	NS	-	-	-

Season	ANOVA Results	Factor Means	
		Fall	Spring
Chironomidae	S	26137.41 (2749.46)	13031.74 (3469.04)
Oligochaeta	S	5626.02 (1496.39)	1541.299 (370.94)
Nematoda	S	5374.40 (1320.65)	1753.43 (541.10)
Ceratapogonidae	NS	-	-
Ephemeroptera (unknown)	NS	-	-

Note: NS = not significant, S = significant. Means with the same letter are not significantly different.

Table 11
Mean Density (Individuals per Square Meter) of Macroinvertebrate Taxa From Pools in
Three Channel Types and Two Sampling Periods

Order	Family	Fall			Spring		
		RC	HC	UC	RC	HC	UC
Bivalvia		3202.25	151.90	374.12	31.65	22.91	16.46
Coleoptera	Dryopidae	0.00	0.00	50.63	0.00	0.00	0.00
	Dytiscidae	0.00	0.00	12.66	0.00	0.00	0.00
	Elmidae	189.03	459.92	232.07	50.63	14.77	141.29
	Gyrinidae	0.00	0.00	0.00	0.00	0.00	0.00
	Staphylinidae	0.00	0.00	12.66	0.00	0.00	0.00
	Unknown	0.00	0.00	0.00	0.00	10.13	0.00
Collembola		25.32	50.63	0.00	16.88	10.13	0.00
Diptera	Ceratopogonidae	354.29	1648.82	874.40	187.76	246.61	265.82
	Chironomidae	25257.38	19656.54	16662.87	10650.21	7309.03	6260.28
	Empididae	46.41	35.44	63.29	0.00	9.21	37.13
	Nymphomyiidae	0.00	0.00	0.00	0.00	31.65	0.00
	Simuliidae	0.00	0.00	0.00	160.34	39.78	20.80
	Tabanidae	0.00	0.00	0.00	0.00	0.00	6.33
	Tanyderidae	0.00	0.00	0.00	0.00	0.00	0.00
	Tipulidae	0.00	0.00	0.00	0.00	0.00	0.00
	Unknown	173.00	0.00	25.32	6.33	12.66	6.75
Ephemeroptera	Baetidae	0.00	0.00	0.00	0.00	18.99	12.66
	Baetiscidae	0.00	0.00	31.65	0.00	0.00	0.00
	Caenidae	0.00	0.00	0.00	84.39	0.00	13.50
	Ephemeridae	75.95	218.81	57.38	0.00	22.78	0.00
	Heptageniidae	482.42	25.32	113.92	49.79	0.00	0.00
	Isonychiidae	75.95	0.00	0.00	0.00	0.00	0.00
	Polmitarcyidae	0.00	0.00	0.00	0.00	0.00	0.00
	Potamanthidae	0.00	0.00	0.00	25.32	10.13	0.00
	Tricorythidae	0.00	25.32	0.00	101.27	0.00	25.32
	Unknown	555.56	510.55	25.32	501.69	120.25	89.03
Gastropoda		1244.73	60.76	215.19	18.99	0.00	0.00
Hirudinea		78.76	0.00	0.00	0.00	0.00	0.00
Hydracarina		1271.31	1475.77	249.65	174.68	424.64	181.56
Isopoda		0.00	0.00	0.00	0.00	16.88	0.00
Megaloptera		0.00	0.00	0.00	0.00	25.32	0.00
Nematoda		6688.47	4557.08	4784.67	1913.92	1646.51	5346.90
Nematomorpha		0.00	0.00	12.66	0.00	0.00	0.00
Odonata	Coenagrionidae	60.34	0.00	0.00	0.00	0.00	20.25
	Corduliidae	0.00	101.27	0.00	0.00	0.00	0.00
	Gomphidae	25.32	33.76	67.51	0.00	10.13	0.00
	Unknown	16.88	25.32	29.54	6.33	0.00	0.00
Oligochaeta		6435.16	2394.09	1088.19	1254.85	752.00	671.91
Orthoptera	Tridactylidae	0.00	0.00	6.33	0.00	25.32	0.00
Plecoptera	Perlidae	168.78	50.63	0.00	56.96	0.00	0.00
	Perlodidae	0.00	0.00	12.66	48.10	23.21	0.00
	Unknown	0.00	37.97	0.00	0.00	14.47	0.00
Trichoptera	Brachycentridae	0.00	0.00	0.00	0.00	0.00	12.66
	Hydropsychidae	126.58	0.00	0.00	0.00	0.00	0.00
	Hydroptilidae	243.32	48.10	75.95	0.00	0.00	25.32
	Leptoceridae	289.73	1185.90	210.97	0.00	0.00	37.13
	Philopotamidae	105.49	0.00	12.66	10.13	0.00	0.00
	Polycentropodidae	12.66	0.00	25.32	0.00	0.00	0.00
	Psychomyiidae	0.00	0.00	0.00	0.00	6.33	0.00
	Unknown	109.70	12.66	50.63	63.29	41.59	40.51

8,000 macroinvertebrates per square meter. In contrast, pools in the historically channelized reach during the present study yielded density estimates of 10,000 to 30,000 individuals per square meter (Table 11). In riffles, density estimates were 2,000 to 12,000 individuals per square meter in 1987-1989 versus 8,000 to 31,000 individuals per square meter (Table 12) in the present study.

The detailed spatial comparison in the recently channelized reach, where two particular pool and riffle pairs were sampled in both the present (post-project) and previous (pre-project) study, did not indicate adverse effects on the macroinvertebrate community. Once again, macroinvertebrate density was substantially greater in the present than previous study. Focusing on fall data, when density was greatest, macroinvertebrate density in riffles averaged 68,000 versus 28,000 individuals per square meter in the present and previous study, respectively. In pools, density of macroinvertebrates in fall equaled 31,000 and 10,000 individuals per square meter in the present and previous studies, respectively. Despite the overall differences in density, community composition was similar in the pre- and post-project studies (Tables 13 and 14). The community was clearly dominated by chironomids in both studies – in both pools and riffles, in both fall and spring. Oligochaetes tended to be the next most abundant group. Nematodes were more abundant in the post-project than pre-project survey.

In both this and the previous study, macroinvertebrate density tended to be greater in fall than spring. Scouring flow in winter and early spring probably accounts for much of the seasonal difference (Payne and Miller 1991). Riffles showed a greater reduction of invertebrate density in spring versus fall in the previous study (Payne et al. 1991; Payne and Miller 1991). In the first study, seasonal differences were more pronounced for fully aquatic oligochaetes than for chironomids with an aerial dispersal adult stage in their life history. In the present study, there was no apparent difference in pool and riffle susceptibility to scour-associated reductions of macroinvertebrate density.

Table 12
Mean Density (Individuals per Square Meter) of Macroinvertebrate Taxa From Riffles in
Three Channel Types and Two Sampling Periods

Order	Family	Fall			Spring		
		RC	HC	UC	RC	HC	UC
Bivalvia		2067.51	87.76	244.73	0.00	16.88	49.91
Coleoptera	Dryopidae	0.00	0.00	0.00	0.00	0.00	0.00
	Dytiscidae	0.00	0.00	0.00	0.00	0.00	0.00
	Elmidae	59.07	50.63	774.26	46.41	6.33	217.00
	Gyrinidae	0.00	0.00	0.00	0.00	0.00	56.96
	Staphylinidae	0.00	0.00	0.00	0.00	14.47	0.00
	Unknown	0.00	35.44	0.00	0.00	0.00	0.00
Collembola		0.00	16.88	0.00	12.66	0.00	0.00
Diptera	Ceratopogonidae	599.16	1504.22	957.81	1069.89	685.53	597.11
	Chironomidae	35240.51	22135.86	21035.86	19333.91	5945.57	13815.73
	Empididae	253.16	322.36	426.16	16.88	46.41	119.53
	Nymphomyiidae	0.00	0.00	0.00	0.00	0.00	0.00
	Simuliidae	0.00	25.32	113.92	240.01	31.65	4028.57
	Tabanidae	0.00	0.00	0.00	0.00	8.44	0.00
	Tanyderidae	0.00	0.00	0.00	0.00	6.33	0.00
	Tipulidae	0.00	59.07	0.00	0.00	6.33	0.00
	Unknown	185.65	23.21	82.28	48.10	82.28	66.00
Ephemeroptera	Baetidae	0.00	166.67	0.00	168.35	52.44	97.65
	Baetiscidae	0.00	0.00	0.00	0.00	0.00	0.00
	Caenidae	25.32	0.00	0.00	26.58	6.33	0.00
	Ephemeridae	0.00	0.00	0.00	0.00	0.00	0.00
	Heptageniidae	550.63	80.17	533.76	195.36	0.00	202.53
	Isonychiidae	0.00	10.13	0.00	0.00	0.00	0.00
	Polmitarcyidae	0.00	16.88	0.00	0.00	0.00	0.00
	Potamanthidae	0.00	0.00	0.00	12.66	0.00	0.00
	Tricorythidae	168.78	0.00	0.00	6.33	0.00	14.47
Unknown	721.52	598.31	1649.79	802.22	341.41	433.09	
Gastropoda		2244.73	109.70	455.70	18.99	0.00	0.00
Hirudinea		0.00	0.00	0.00	329.11	0.00	0.00
Hydracarina		1500.00	1677.22	6713.08	442.74	349.91	899.52
Isopoda		194.09	0.00	0.00	67.51	0.00	6.33
Megaloptera	Sialidae	0.00	0.00	0.00	0.00	0.00	0.00
Nematoda		10219.41	1528.27	4375.53	2001.63	458.05	2800.60
Nematomorpha		0.00	0.00	25.32	0.00	0.00	0.00
Odonata	Coenagrionidae	0.00	0.00	25.32	0.00	0.00	0.00
	Corduliidae	0.00	0.00	0.00	0.00	0.00	0.00
	Gomphidae	0.00	0.00	0.00	0.00	0.00	0.00
	Unknown	0.00	0.00	0.00	0.00	0.00	0.00
Oligochaeta		10841.77	1122.78	4913.50	1709.55	1159.19	1755.15
Orthoptera	Tridactylidae	0.00	0.00	0.00	0.00	12.66	0.00
Plecoptera	Perlidae	375.53	0.00	270.04	0.00	0.00	0.00
	Perlodidae	6.33	33.76	29.54	83.12	6.33	73.24
	Unknown	25.32	45.57	75.95	40.08	0.00	0.00
Trichoptera	Brachycentridae	0.00	0.00	25.32	33.76	0.00	0.00
	Hydropsychidae	12.66	435.44	276.37	16.88	12.66	113.92
	Hydroptilidae	497.89	97.05	824.89	16.88	12.66	45.21
	Leptoceridae	721.52	110.97	582.28	25.32	23.21	63.29
	Philopotamidae	455.70	77.22	2318.57	18.99	0.00	516.58
	Polycentropodidae	101.27	6.33	0.00	0.00	0.00	0.00
	Psychomyiidae	0.00	10.13	0.00	0.00	0.00	0.00
Unknown	565.40	510.13	907.17	130.38	84.09	450.27	

Table 13
Relative Abundance (Percent) of Macroinvertebrate Taxa From Pools
in the Recently Altered Portion of Luxapalila Creek Among Pre- and
Post-Alteration Sampling Periods and Two Seasons

Invertebrate Group	Fall		Spring	
	Pre	Post	Pre	Post
Amphipoda	0.15	0.00	0.28	0.00
Bivalvia	9.44	6.79	5.50	0.21
Coleoptera	0.53	0.40	0.46	0.33
Collembola	0.23	0.05	0.37	0.11
Diptera – Ceratopogonidae	1.90	0.75	1.38	1.22
Diptera – Chironomidae	47.91	53.58	60.55	69.13
Diptera – Empididae	0.08	0.10	0.00	0.00
Diptera – Simuliidae	0.00	0.00	0.00	1.04
Diptera – Tanyderidae	0.08	0.00	0.00	0.00
Ephemeroptera	0.15	2.52	0.09	4.95
Gastropoda	0.46	2.64	0.83	0.12
Heteroptera	0.00	0.00	0.00	0.00
Hirudinea	0.23	0.17	0.18	0.00
Hydracarina	3.88	2.70	1.65	1.13
Isopoda	0.23	0.00	0.18	0.00
Nematoda	7.31	14.19	6.15	12.42
Nemertea	0.46	0.00	0.09	0.00
Odonata	0.30	0.22	0.09	0.04
Oligochaeta	23.69	13.65	20.00	8.14
Platyhelminthes	0.99	0.00	1.01	0.00
Plecoptera	0.08	0.36	0.37	0.68
Polychaeta	0.08	0.00	0.00	0.00
Trichoptera	1.83	1.88	0.83	0.48

Table 14
Relative Abundance (Percent) of Macroinvertebrate Taxa From
Riffles in the Recently Altered Portion of Luxapalila Creek Among
Pre- and Post-Alteration Sampling Periods and Two Seasons

Invertebrate Group	Fall		Spring	
	Pre	Post	Pre	Post
Amphipoda	0.00	0.00	0.10	0.00
Bivalvia	6.08	3.07	5.93	0.00
Coleoptera	0.14	0.09	0.15	0.17
Collembola	0.31	0.00	0.10	0.05
Diptera – Ceratopogonidae	0.08	0.89	0.00	3.98
Diptera – Chironomidae	53.00	52.25	46.76	71.96
Diptera – Empididae	0.02	0.38	0.00	0.06
Diptera – Simuliidae	0.00	0.00	1.89	0.89
Diptera – Tanyderidae	0.00	0.00	0.00	0.00
Ephemeroptera	0.88	2.17	2.69	4.51
Gastropoda	1.33	3.33	0.45	0.07
Heteroptera	0.00	0.00	0.05	0.00
Hirudinea	0.00	0.00	0.05	1.23
Hydracarina	18.97	2.22	7.08	1.65
Isopoda	0.00	0.29	0.00	0.25
Nematoda	0.88	15.15	1.60	7.45
Nemertea	0.74	0.00	0.30	0.00
Odonata	0.05	0.00	0.05	0.00
Oligochaeta	10.17	16.07	19.74	6.36
Platyhelminthes	0.71	0.00	1.99	0.00
Plecoptera	0.24	0.60	0.35	0.46
Polychaeta	0.00	0.00	0.00	0.00
Trichoptera	6.39	3.49	10.72	0.90

4 Summary

Water and substratum characteristics are generally similar for riffles or pools whether in the downstream and recently channelized reach, the middle reach that has not been channelized, or the upper reach that was straightened and simplified in 1922. The sandy and unsorted gravelly substratum of pools and riffles, respectively, along with good water quality throughout, largely determines the macroinvertebrate community of this stream. Functionally, the macroinvertebrate community was similar throughout the stream, regardless of channel type or if samples had been from riffles or pools. A possible exception was the greater abundance of filter-feeding *C. fluminea* in the downstream, recently channelized reach. However, this might have had more to do with stream trophic dynamics rather than channel modifications.

Small, short-lived animals such as chironomid larvae, oligochaetes, nematodes, and the Asian clam, *C. fluminea*, dominated the stream macroinvertebrate community. The density of these invertebrates tended to increase in a downstream direction; this trend probably reflected the relatively oligotrophic nature of the upstream reaches of this clearwater stream. The canopy-less nature of the recently channelized downstream reach probably allows greater levels of autochthonous production of organic carbon during low flow than is possible in the relatively heavily shaded, narrower middle and upper reaches. Regardless of cause, it is clear that secondary production, reflected by macroinvertebrate density in general, and *C. fluminea* density in particular, increases moving downstream.

Overall, general limnological and trophic considerations appeared to overwhelm spatial patterns in biological organization that might otherwise have promoted differences among recently channelized, unchannelized, and historically channelized reaches. Dominance of small macroinvertebrates, most of which can rapidly recolonize disturbed areas, encouraged similarity rather than difference in community composition of the three types of stream reach. Indeed, even pools and riffles in this stream were similar. Pools were poorly formed and small, only slightly deeper than riffles, and probably subjected to considerable scour during high discharge. This was probably more true after than before project construction in the recently channelized reach. Gravel in this sand-channel stream was mostly limited to riffles, poorly sorted (Payne et al. 1991), and probably unstable during very high flows.

Almost certainly, relatively long-lived and sedentary animals, such as native unionid mussels, are more likely to thrive in the unchannelized than recently

channelized reaches of the stream, and even in that reach truly stable shoals of sand and gravel barely exist. Compared with *C. fluminea*, most native unionids require long-term stability of sand or gravel substratum to establish moderately dense assemblages (Payne et al. 1989). Long-term monitoring will be necessary to evaluate the status of the sparse unionid fauna in the unchannelized middle reach of the river.

The ability of the Luxapalila Creek macroinvertebrate fauna to recolonize disturbed areas was evident from the pre- and post-project comparisons of the recently channelized lower reach of the stream. Short-lived and relatively small chironomids, oligochaetes, nematodes, and *C. fluminea*, that in combination comprise the very large majority of the macroinvertebrate fauna, fully recolonized the recently channelized lower reach. Both in that reach in Columbus, MS, and in the historically channelized but not recently disturbed reach in Alabama, basic community composition was the same as in 1987-1989. In addition, total density of macroinvertebrates was higher in the present study. There was no evidence of the macroinvertebrate community of Luxapalila Creek having been detrimentally affected by flood control measures. Indeed, production of dominant organisms probably increased.

References

- Barbour, M. T., Gerritsen, J., Snyder, B. D., and Stribling, J. B. (1999). *Rapid bioassessment protocols for use in streams and wadeable rivers: Periphyton, benthic macroinvertebrates and fish*. 2nd ed., EPA 841-B-99-002, U.S. Environmental Protection Agency, Office of Water, Washington, DC.
- Jandel Corporation. (1995). "SigmaStat for Windows version 2.0 statistical software," San Rafael, CA.
- Miller, A. C., and Bingham, C. R. (1987). "A hand-held benthic core sampler," *Journal of Freshwater Ecology* 4: 77-81.
- Payne, B. S., and Miller, A. C. (1991). "The structure of dominant invertebrate assemblages in a small southeastern stream," *Journal of Freshwater Ecology* 6: 257-266.
- Payne, B. S., Miller, A. C., Hartfield, P. D., and McMahon, R. F. (1989). "Variation in size demography of lotic populations of *Corbicula fluminea* (Muller)," *The Nautilus* 103: 78-82.
- Payne, B. S., Miller, A. C., Miller-Way, C., and Bingham, C. R. (1991). "Macroinvertebrates of Luxapalila Creek, Mississippi and Alabama, 1987-89," Miscellaneous Paper EL-91-5, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Standford, J. A., and Ward, J. V. (1979). "Stream regulation in North America." *The ecology of regulated streams*. J. V. Ward and J. A. Stanford, ed., Plenum Press, New York, 215-236.
- Zar, J. H. (1987). *Biostatistical analysis*. 2nd ed., Prentice Hall, Englewood Cliffs, NJ.

